



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Unidirectional Na⁺ Influx under Different Growth Stages of *Puccinellia tenuiflora* and Wheat

C. M. Wang
Lanzhou University, China

J. L. Zhang
Lanzhou University, China

G. Q. Wu
Lanzhou University, China

Suomin Wang
Lanzhou University, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/2-2/44>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Unidirectional Na^+ influx under different growth stages of *Puccinellia tenuiflora* and wheat

C.M. Wang, J.L. Zhang, G.Q. Wu, S.M. Wang*

School of Pastoral Agriculture Science and Technology, Lanzhou University, Lanzhou, Gansu, 730000,

* E-mail: smwang@lzu.edu.cn

Key words: isotope ($^{22}\text{Na}^+$) tracing technique, unidirectional Na^+ influx, growth stage, halophyte, glycophyte

Introduction Restricting unidirectional Na^+ influx into root was an important way to reduce the net accumulation of Na^+ . However, there was no report regarding whether the unidirectional Na^+ influx was affected by the growth stage. Consequently, it was hard to compare the influxes between/among different growth stages or species. This study was focused on this problem. Thus, Gramineous halophyte *P. tenuiflora* and glycophyte wheat were chosen, for testing the unidirectional Na^+ influxes of seedlings at the 2 leaf, 3 leaf and 4 leaf stages in 25 and 100 mM NaCl treatments.

Materials and methods Seedlings of *P. tenuiflora* and wheat were germinated and then cultured in modified Hoagland's nutrient solution, and subsequently NaCl treated for 7 days after reaching the 2, 3 and 4 leaf stage. Seedlings were harvested and weighed. $^{22}\text{Na}^+$ influx was evaluated according to the method described by Essah et al. (2003).

Results With advancing plant growth (Figure 1), unidirectional Na^+ influx was increased significantly in both species, and more obvious in 100 mM NaCl (Figure 2). There was no significant difference between *P. tenuiflora* and wheat in 25 mM NaCl. But in 100 mM NaCl, *P. tenuiflora* had stronger ability than wheat to restrict the increase of unidirectional Na^+ influx in the 4 leaves stage.




Figure 1 Root fresh weight of *P. tenuiflora* and wheat at different growth stages ($P < 0.05$, Duncan test).




Figure 2 Unidirectional Na^+ influx of *P. tenuiflora* and wheat at different growth stages treated with 25 and 100 mM NaCl.

Conclusions Unidirectional Na^+ influx was increased significantly with growth stage in both species, but more obvious in 100 mM NaCl. At the 2-3 leaf stage, the capacities of restricting Na^+ influx in both *P. tenuiflora* and wheat were similar, but significantly enhanced in *P. tenuiflora* after more mature (4 leaves). It was recommended that the growth stages should be taken into account when unidirectional Na^+ influxes were compared between/among different growth stages or species.

References

- Tester M, Davenport R (2003). Na^+ tolerance and Na^+ transport in higher plants. *Annals of Botany* 91: 1-25.
 Essah PA, Davenport R, Tester M (2003). Sodium influx and accumulation in *Arabidopsis*. *Plant Physiology* 133: 307-318.